

THE UNITED SHAYES OF AMERICA

To all to whom these presents shall come; Hinneer Hi-Bred International, Inc.

THEOLOGIS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY VIALS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE UGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR PORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT TO BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PHAR1'

In Vestimonn Marrest, I have hereunto set my hand and caused the seal of the Plant Anxiety Protection Office to be affixed at the City of Washington, D.C. this twenty-third day of November, in the year two thousand and soven.

Au.

ger zun

Commissioner Plant Variety Protection Office Agricultural Marketing Service Secretary

REPRODUCE LOCALLY. Include form number and	date on all repro	ductions		Form Approved - OMB No. 0581-0055
IJ.S. DEPARTMEI AGRICULTURAL I SCIENCE AND TECHNOLOGY - P	MARKETING SER	VICE	The following statements are made in a the Paperwork Reduction Act (PRA) of	ocordance with the Privacy Act of 1974 (5 U.S.C. 552a) and 1995.
APPLICATION FOR PLANT VA (Instructions and information co				nine if a plant variety protection certificate is to be issued nfidential until certificate is issued (7 U.S.C. 2426).
. NAME OF OWNER	·····	· · · · · · · · · · · · · · · · · · ·	2, TEMPORARY DESIGNATION OR EXPERIMENTAL NAME	3. VARIETY NAME
Pioneer Hi-Bred	Internatio	nal, Inc.		PHAR1
ADDRESS (Street and No., or R.F.D. No., City,	State, and ZIP Co	de, and Country)	5. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY
7204 NW	62 nd Avenı		515/270-4051	PVPO NUMBER
	, IA 50131-		6. FAX (include area code)	9 000 000 000
oomioton,	, 1, 1, 00 101		515/253-2125	200500253 FILING DATE
, IF THE OWNER NAMED IS NOT A "PERSON",	GIVE FORM OF	8. IF INCORPORATED, GIVE	9. DATE OF INCORPORATION	
ORGANIZATION (corporation, partnership, asso Corporation	ciation, etc.)	STATE OF INCORPORATION	March 5, 1999	May 17,2005
0. NAME AND ADDRESS OF OWNER REPRES	ENTATIVE(S) TO	SERVE IN THIS APPLICATION. (First p.	person listed will receive all papers)	F FILING AND EXAMINATION FEES: 8 3652.00
<u>_</u>		R. Anderson		R DATE 5/17/05 C CERTIFICATION FEE:
Rese		Product Development), Box 85		\$ 768.00
		, IA 50131-0085		E DATE 10/15/07
				D DATE 10/13/-1
TELEPHONE (Include area code)	12. FAX (Includ	le area code)	13. E-MAIL	
515/270-4051		515/253-2125	stev	en.anderson@pioneer.com
4. CROP KIND (Common Name)	16. FAMILY N	AME (Botanical)	18. DOES THE VARIETY CONTA	IN ANY TRANSGENES? (<i>OPTIONAL</i>)
5. GENUS AND SPECIES NAME OF CROP	17. IS THE VA	Gramineae RIETY A FIRST GENERATION HYBRIC	☐ YES ☑ NO	COLONED LINE A ADUIN DEFENDANCE MUNICIPE SOO TUE
Zea Mays	☐ YES	⊠ NO	APPROVED PETITION TO D COMMERICALIZATION.	SSIGNED USDA-APHIS REFERENCE NUMBER FOR THE EREGULATE THE GENETICALLY MODIFIED PLANT FOR
 CHECK APPROPRIATE BOX FOR EACH ATT (Follow instructions on reverse) 		ITTED		Y THAT SEED OF THIS VARIETY BE SOLD AS A CLASS Section 83(a) of the Plant Variety Protection Act)
 a.			☐ YES (If "yes", answer 21. DOES THE OWNER SPECIF' NUMBER OF CLASSES?	items 21 and 22 below) 🔲 NO (if "no", go to item 23) Y THAT SEED OF THIS VARIETY BE LIMITED AS TO
d, X Exhibit D. Additional Description of the V e. ⊠ Exhibit E. Statement of the Basis of the G f. ⊠ Voucher Sample (2,500 viable untreated)	Owner's Ownership		☐ YES ☐ NO	
verification that tissue culture will be deper repository) g. ⊠ Filing and Examination Fee (\$3,652), ma	osited and maintain de payable to "Tre	ned in an approved public		☐ FOUNDATION ☐ REGISTERED ☐ CERTIFIED Y THAT SEED OF THIS VARIETY BE LIMITED AS TO 17
States" (Mail to the Plant Variety Protecti	on Office)			
			IF YES, SPECIFY THE NUMB	ER 1,2,3, etc. FOR EACH CLASS.
			FOUNDATION RE	GISTERED CERTIFIED
				cessary, please use the space indicated on the reverse.)
3. HAS THE VARIETY (INCLUDING ANY HARVE FROM THIS VARIETY BEEN SOLD, DISPOSE OTHER COUNTRIES?			24. IS THE VARIETY OR ANY CO INTELLECTUAL PROPERTY	MPONENT OF THE VARIETY PROTECTED BY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)?
☑ YES ☐ NO			☐ YES ☑ NO	
IF YES, YOU MUST PROVIDE THE DATE OF FOR EACH COUNTRY AND THE CIRCUMSTA	FIRST SALE, DISI NCES. <i>(Please u</i>	POSITION, TRANSFER, OR USE se space indicated on reverse.)	IF YES, PLEASE GIVE COUN' REFERENCE NUMBER. (Plea	TRY, DATE OF FILING OR ISSUANCE AND ASSIGNED ase use space indicated on reverse.)
15. The owners declare that a viable sample of ba for a tuber propagated variety a tissue culture	sic seed of the var will be deposited in	iety has been furnished with application n a public repository and maintained fo	n and will be replenished upon request in a r the duration of the certificate.	accordance with such regulations as may be applicable, or
The undersigned owner(s) is(are) the owner of entitled to protection under the provisions of Se	this sexually repro ction 42 of the Pla	duced or tuber propagated plant variety int Variety Protection Act.	/, and believe(s) that the variety is new, di	stinct, uniform, and stable as required in Section 42, and is
Owner(s) is (are) informed that false representa	ition herein can je	opardize protection and result in penalti	es.	
IIGNATURE OF OWNER	1 111 11 11 11 11 11	S	IGNATURE OF OWNER Attack	en & Anderson
IAME (Please print or type)		N	AME (Please print or type)	v v pur vice
		,	Steve	en R. Anderson
APACITY OR TITLE	DATE	. C	APACITY OR TITLE	DATE
			Research Scientist	5-16-2005

application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that vill reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 illing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials o make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuanc of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvpindex.htm

Fo avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

TEM

19a.Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- I9c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

United States, Nov. 1, 2004; Japan Nov. 1, 2004; France, Nov. 1, 2004

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the rariety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing astructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

he U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, olitical beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information 3 raille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

o file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD) SDA is an equal opportunity provider and employer.

iT-470 (04-03) designed by the Plant Variety Protection Office using Word 2002.

Exhibit A: Developmental history for PHAR1

Pedigree: PH2VJ/PH3RC)X7231221X

Pioneer Line PHAR1, Zea mays L., a yellow endosperm, dent corn, inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross hybrid PH2VJ (PVP Certificate Number 9900364) X PH3RC (PVP Certificate Number 200200257) using the pedigree method of plant breeding. Varieties PH2VJ and PH3RC are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing was practiced from the above hybrid for 9 generations using pedigree selection. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Macomb, Illinois as well as other Pioneer research locations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations again made for uniformity.

Variety PHAR1 has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 7 generations with careful attention paid to selection criteria and uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity and stability, and for 3 generations during the final stages of inbred development and seed multiplication. Very high standards for genetic purity have been established morphologically using field observations and using sound lab molecular marker methodology.

No variant traits have been observed or are expected in PHAR1.

The criteria used in the selection of PHAR1 were yield, both per se and in hybrid combinations. Late season plant health and late season root lodging, grain quality, stalk lodging resistance, and kernel size are especially important in production, and were also important criteria considered during selection.

Exhibit A: Developmental history for PHAR1

Pedigree Grown	Inbreeding Level of
Season/Year	Pedigree Grown
PH2VJ	F0
Fall 1996	
PH3RC	F0
Fall 1996	
PH2VJ/PH3RC	F1
Winter 1996	
PH2VJ/PH3RC)X	F2
Summer 1997	
PH2VJ/PH3RC)X7	F3
Summer 1998	
PH2VJ/PH3RC)X72	F4
Winter 1998	
PH2VJ/PH3RC)X723	F5
Summer 1999	
PH2VJ/PH3RC)X7231	F6
Winter 1999	
PH2VJ/PH3RC)X72312	F7
Summer 2000	
PH2VJ/PH3RC)X723122	F8
Winter 2000	
PH2VJ/PH3RC)X7231221	F9
Summer 2001	
PH2VJ/PH3RC)X7231221X	F10 (Seed)

^{*}PHAR1 was selfed and ear-rowed from F3 through F9 generation.
#Uniformity and stability were established from F7 through F9 generation and beyond when seed supplies were increased.

Variety PHAR1 mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PH2VJ (PVP Certificate No. 9900364). Table 1 shows two sample t-tests on data collected primarily in Johnston and Dallas Center, Iowa in 2003 and 2004. The traits collectively show measurable differences between the two varieties.

Exhibit B: Novelty Statement

Variety PHAR1 has shorter shank length (7.1 cm vs 11.0 cm) than variety PH2VJ (Table 1).

Variety PHAR1 has shorter ear length (14.1 cm vs 15.8 cm) than variety PH2VJ (Table 1).

Variety PHAR1 has a light green silk color (Munsell 10Y 7/6) and PH2VJ has a pink silk color (Munsell 2.5R 6/6).

Silk color establishes a clear difference between PHAR1 and PH2VJ.

Exhibit B: Novelty Statement Table(s)

Table 1: Data from Johnston and Dallas Center, Iowa in 2003 and 2004 presented by trait, across years, and broken out by year. Data are supporting evidence for differences between PHAR1 and PH2VJ. Each year varieties were grown in 3 locations that had different environmental conditions. Environments had different planting dates and were in different fields. A two-sample t-test was used to compare differences between means.

Shank length (cm)											٠					
Level Over All	Station	Year	Variety-1 PHAR1	Variety-2 PH2VJ	Cnt-1 30	Cnt-2 30	Mean-1 7.1	Меап-2 11.0	Mean_Diff -3.9	StDev-1 1.447	StDev-2 2.484	StErr-1 0.264	StErr-2 0.454	DF 58	t-Value -7.5	Prob_Pool 0.000
Year		2003	PHARI	PH2VJ	15	15	7.8	12.8	-5.0	1.373	1.859	0.355	0.480	28	-8.4	0.000
Year		2004	PHAR1	PH2VJ	15	15	6.4	9.3	-2.9	1.183	1.624	0.306	0.419	28	-5.5	0.000
Ear length (cm)		ļ														
Level	Station	Year		Variety-2	Cnt-1	Cnt-2	Mean-1	Mean-2	Mean_Diff	StDev-1	StDev-2	StErr-1	StErr-2	DF	t-Value	Prob_Pool
Over All			PHAR1	PH2VJ	30	30	14.1	15.8	-1.7	0.937	1.031	0.171	0.188	58	-6.6	0.000
Year		2003	PHAR1	PH2VJ	15	15	14.0	16.0	-2.0	1.134	1.309	0.293	0.338	28	-4.5	0.000
Year		2004	PHAR1	PH2VJ	15	15	14.3	15.6	-1.3	0.704	0.632	0.182	0.163	28	-5.5	0.000

Based on previous discussions with the PVP office the traits longitudinal creases and marginal leaf waves were not collected. These traits have low distinguishing power and are variable due to daily fluctuations in water status of the plants. Therefore, we eliminated them from our process based on previous feedback from the PVP office. For insect or disease traits we included data from disease pressure locations only if they were available and paired with the public check. Most often diseases and insect trials are conducted on hybrids since that is the product ultimately sold. In addition, creating consistent disease pressure and infestation levels is costly and difficult.

In cases where less than 15 observations are presented the trait was collected at the plot level as it always has been done in the past. This means many more plants were visually evaluated according to the procedure outlined below, and then a score of the "population" of the plants was recorded for each location.

The experimental design and methods for 2004 were as follows:

The experiment procedures involved three environments with different planting dates per year, planted in 17.42 ft. rows with 2 rows for each variety. Approximately 24-30 plants emerged in each of 2 rows for a total of around 48 to 60 plants being evaluated at each location and 144 to 180 plants across locations. For plant level traits, we sampled 5 representative plants from the 2 rows of the 2 row plot (group) of plants at each location. For plot level traits we evaluated the 2 row plot (group) and gave a representative score or average on the 48-60 plants in the group within an experiment.

Some traits can be especially variable under different environmental factors influenced by weather, soil type, or planting dates. Varying temperatures or day length could impact the meristem growth during various tissue differentiation stages. The meristem differentiation of the ear and other tissues could be impacted as well as the success of pollination during flowering and frequency of kernel abortion during grain fill.

We have included weather data in the table that follows.

	300	מעשכ מע	A CHINETED	7000	Ē	T. H. C.L.	I.	
Month		CINCHING DEGREE UNITS (GDUS)	02100	(8,000	Ξ.	PKECIFII ALIO	ION (Inches	(S)
	20	2003	20	2004	20	2003	20	2004
	D. Center	Johnston	D. Center	Johnston		3. Center Johnston	D. Center	Johnston
May	375	380	548	527		5.43	7 19	7.75
June	909	604	609	610	192	4 23	1 97	900 6
July	628	782	723	736	0.18	3.4	2 20	200
August	795	786	612	615	0 44	0.51	1 95	10.1
September	456	468	598	560	2.19	2.52	138	5 7
TOTAL	2860	3020	3090	3048	10.43	16.09	14.78	24 87

Calculate GDU's
Growing Degree Units use following formula: GDU = ((T1+T2)/2)-50
Where T1 = minimum temperature for a given day with 50 degrees Fahrenheit as the minimum temperature used and 86 degrees Fahrenheit is the maximum temperature used.
Where T2 = maximum temperature for a given day with 86 degrees Fahrenheit as the maximum temperature used and 50 degrees Fahrenheit is the minimum temperature used.

United States Department of Agriculture, Agricultural Marketing Service Science and Technology, Plant Variety Protection Office National Agricultural Library Building, Room 400 Beltsville, MD 20705-2351

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Pioneer Hi-Bred Int		l vallety Seed S	source		HAR1	emporary De	esignation
	o., or R.F.D. No., City, State, nue, P.O. Box 85, Johnston,		i FOR OF	FICIAL USE	1	PVPO Nu 0 0 5	mber 0 0 2 5
adding leading zeroe	e number that describes the vestif necessary. Completeness y for an adequate variety des	should be striven for to	establish an adequa	y in the spac ate variety de	es below. Right j escription. Traits o	ustify whole designated b	numbers by by a "*" are
COLOR CHOICES (I 01. Light Green 02. Medium Green 03. Dark Green 04. Very Dark Gree 05. Green-Yellow	Use in conjunction with Munso 06. Pale Yellow 07. Yellow 08. Yellow-Orange n 09. Salmon 10. Pink-Orange	ell color code to describe 11. Pink 12. Light Red 13. Cherry Red 14. Red 15. Red & White	all color choices; de 16. Pale Purple 17. Purple 18. Colorless 19. White 20. White Cappe	21. 22. 23. 24.	and #26 in Comm Buff Tan Brown Bronze Variegated (Desc	26. Other (I	
Yellow Dent Familie Family B14 B37 B73 C103 Oh43	O CHOICES [Use the most sires: Members CM105, A632, B64, B68 B37, B76, H84 N192, A679, B73, Nc268 Mo17, Va102, Va35, A682 A619, MS71, H99, Va26 W64A, A554, A654, Pa91	nilar (in background and i Yellow Dent (Unrelated) Co109, ND246 Oh7, T232 W117, W153R W182BN White Dent: Cl66, H105, Ky2	:	Swe Pop	eet Corn: C13, Iowa512: corn:	5, P39, 2132 722, HP301,	2 HP7211
	intermediate types in "Commo it, 2=Dent, 3=Flint, 4=Flour, 5 t		Pipecorn)	}	Standard Inbred N 2 Type	Name	B73
	DEVELOPED IN THE U.S.A st, 2=N.Central, 3=N.East, 4=		S.West, 7=Other		Standard Seed Se Region	ource	PI 550473
	egion Best Adaptability; show HEAT UNITS 1,395.5 From emergence 1,394.7 From emergence 52 From 10% to 90% From 50% silk to From 50% silk to	to 50% of plants in silk to 50% of plants in poller	·	 	DAYS 65 64 2	<u>1</u> .	JNITS 382.5 366.8 49
82.2 cm Ear I 15.5 cm Leng 0.0 Average 1.0 Average	Height (to tassel tip) Height (to base of top ear nod th of Top Ear Internode Number of Tillers Number of Ears per Stalk anin of Brace Roots: 1=Abser	e) nt, 2=Faint, 3=Moderate,	15.46 8.62 2.30 0.01 0.05 4=Dark	Size I 30 I 30 I 30 I 6 I 6 I	Mean 238.2 98.1 15.5 0.0 1.0 2	St Dev. 19.92 16.92 1.68 0.02 0.07	Sample Size 30 30 30 6 6
Application Variety L	Data	F	Page 1	1 8	Standard Inbred D	Data	

5. LEAF					
	St.Dev.	Sample Size I	Mean	St.Dev.	Sample Siz
10.0 cm Width of Ear Node Leaf	0.98	30 I	Mean 8.7	0.79	
82.2 cm Length of Ear Node Leaf	3.34	30 I	80.9	3.32	<u>3</u> 3
6.6 Number of leaves above top ear	<u>3.34</u> 0.67	30 i 30 i			2
24.0 Degrees Leaf Angle			6.4	<u>0.77</u>	<u>3</u>
	<u>5.92</u>	<u>30</u> !	<u>22.4</u>	<u>5.08</u>	<u> </u>
(Measure from 2nd leaf above ear at anthesis to stalk abov	ve lear)	!	4 (8.6	0-4-1-2-56	a) cole
4 Leaf Color (Munsell Code) 5GY34	O 111	·		Code) <u>7.50</u>	<u> 5 Y 3/4</u>
5 Leaf Sheath Pubescence (Rate on scale from 1=none to	9=like peach i	uzz) i	<u>3</u>		
Marginal Waves (Rate on scale from 1=none to 9=many)	and)	1			
Longitudinal Creases (Rate on scale from 1=none to 9=m	апу)	'	 .		
TASSEL:	St.Dev.	Sample Size I	Mean	St.Dev.	Sample Siz
4.3 Number of Primary Lateral Branches	<u>1.53</u>	<u>30</u> l	<u>7.3</u>	1.30	3
39.8 Degrees Branch Angle from Central Spike	19.41	<u>30</u> 1	15.6	8.41	3
55.1 cm tassel Length	4.16	30 I	55.4	4.16	3
(from top leaf collar to tassel tip)		i			
4 Pollen Shed (Rate on scale from 0=male sterile to 9=hear	vv shed)	i	5		
6 Anther Color (Munsell Code) 5Y76	-,,	i	7 (Munsell	Code) <u>5Y8</u>	54
2 Glume Color (Munsell Code) 5GY56		i		Code) 5GY	
1 Bar Glumes (Glume Bands): 1=Absent, 2=Present		j	1		7-
ı. EAR (Unhusked Data):		, 1			1
Silk Color (3 days after emergence) (Munsell Code)	<u>10`</u>			Code <u>2.50</u>	
2 Fresh Husk Color (25 days after 50% silking) (Munsell Color)		<u>Y6∤8</u> , I		Code <u>5GY</u>	
21 Dry Husk Color (65 days after 50% silking) (Munsell Cod		<u>Y8.54</u> I	21 Munsell	Code <u>2.5</u> Y	<u> (8.54</u>
2 Position of Ear at Dry Husk Stage: 1=Upright, 2=Horizon		t i	<u>3</u>		•
5 Husk Tightness (Rate on scale from 1=very loose to 9=very)	ery tight	. [<u>න</u> න න		
2 Husk Extension (at harvest): 1=Short(ears exposed), 2=N	vledium (<8cm	i), 3=Long 1	<u>3</u>		
(8-10cm beyond ear tip), 4=Very Long (>10cm)		i			
o. EAR (Husked Ear Data)	St. Dev.	Sample Size I	Mean	St.Dev.	
14.2 cm Ear Length	OL DEV.	ו שאנט שועווווגע			Sample Siz
	1 01				•
	<u>1.01</u>	<u>30</u> I	<u>13.1</u>	<u>1.09</u>	· <u>3</u>
43.1 mm Ear Diameter at mid-point	<u>1.95</u>	30 I 30 I	<u>13.1</u> <u>44.6</u>	<u>1.09</u> <u>3.23</u>	. 23
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight	1.95 23.20	30 30 30	<u>13.1</u> <u>44.6</u> <u>118.8</u>	1.09 3.23 43.50	· 33
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows	<u>1.95</u>	30 I 30 I	13.1 44.6 118.8 17.9	<u>1.09</u> <u>3.23</u>	. 3333
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct	1.95 23.20 1.01	30 30 30 30 30	<u>13.1</u> <u>44.6</u> <u>118.8</u>	1.09 3.23 43.50	· 33
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly-Curved, 3=Spiral—	1.95 23.20 1.01	30 30 30 30 1	13.1 44.6 118.8 17.9 2	1.09 3.23 43.50 1.74	3333
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 7.1 cm Shank Length	1.95 23.20 1.01	30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4	1.09 3.23 43.50	Sample Siz
 43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly-Curved, 3=Spiral 	1.95 23.20 1.01	30 30 30 30 1	13.1 44.6 118.8 17.9 2	1.09 3.23 43.50 1.74	Sleafest features
mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extree KERNEL (Dried):	1.95 23.20 1.01 1.45 eme conical St.Dev.	30 30 30 30 1 30 30 1 Sample Size	13.1 44.6 118.8 17.9 2 2 7.4 2	1.09 3.23 43.50 1.74 2.11	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral—cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral— 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral—7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Thickness	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral— 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral—7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (Shape Grade)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (describe	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e)	Sample Si
A3.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extremation 10.4 mm Kernel Length 1.7 mm Kernel Width 1.7 mm Kernel Width 1.7 mm Kernel Thickness 1.6 mm Kernel Thickness 1.6 mm Kernel Shape Grade 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating of 1 Aleurone Color (Munsell Code) 1.04 mm Kernel Thickness 1.04 mm Kernel Thick	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (describe 7 Munsell	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2.	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral- 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (1) 7 Aleurone Color (Munsell Code) 1 10YR8/14	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1: 1 (describe 7 Munsell 7 Munsell	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (Pattern Color (Munsell Code) 10 (10 (10 (10 (10 (10 (10 (10 (10 (10	1.95 23.20 1.01 1.01 St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (describe 7 Munsell	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (12 Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 10YR861	1.95 23.20 1.01 1.01 St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1: 1 (describe 7 Munsell 7 Munsell	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral— 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (7 Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High I	1.95 23.20 1.01 1.01 St.Dev. 1.10 0.83 0.86 11.52 (describe)	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1: 1 (describe 7 Munsell 7 Munsell	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral— 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating of Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 1 Sendosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High I (se), 9=High Oil, 10=Other 29.8 gm Weight per 100 kernels (unsized sample)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe) 4 2 Normal Starc Lysine, 8=Sup	30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (described of the second of the	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code E) Code	Sample Si 5Y8/12 0YR7/12
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly-Curved, 3=Spiral—7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (ERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating, 7 Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 1 Sendosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High I (se), 9=High Oil, 10=Other 29.8 gm Weight per 100 kernels (unsized sample)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe) 4 2 Normal Starc Lysine, 8=Sup	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (described of the second of the	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10 e) 7.63	Sample Size
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral—7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre (KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (7 Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 2 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High (se), 9=High Oil, 10=Other 29.8 gm Weight per 100 kernels (unsized sample)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe) 4 2 Normal Starc Lysine, 8=Sup	30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (described of the second of the	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10 e) 7.63	Sample Si
43.1 mm Ear Diameter at mid-point 119.1 gm Ear Weight 15.5 Number of Kernel Rows 2 Kernel Rows: 1=Indistinct, 2=Distinct 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral— 7.1 cm Shank Length 1 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extre KERNEL (Dried): 10.4 mm Kernel Length 7.7 mm Kernel Width 5.6 mm Kernel Width 5.6 mm Kernel Thickness 68.5 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating, 7 Aleurone Color (Munsell Code) 7 Hard Endosperm Color (Munsell Code) 1 Sendosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High I (se), 9=High Oil, 10=Other 29.8 gm Weight per 100 kernels (unsized sample)	1.95 23.20 1.01 1.45 eme conical St.Dev. 1.10 0.83 0.86 11.52 (describe) 4 2 Normal Starc Lysine, 8=Sup	30 30 30 30 30 30 30 30	13.1 44.6 118.8 17.9 2 2 7.4 2 Mean 11.0 6.5 4.1 10.1 1 (described of the second of the	1.09 3.23 43.50 1.74 2.11 St.Dev. 1.61 0.68 0.48 5.29 e) Code 2 Code 10 e) 7.63	Sample Size

Note: Use chart on first page to choose color codes for color traits

Standard Inbred Data

Southern Leaf Blight (Bipolaris maydis) Southern Rust (Puccinia Polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi Other (Specify) C. Stalk Rots	Race Race Race		Anthracnose Leaf Blight Common Rust Common Smut Eyespot Goss's Wilt 3 Gray Leaf Spot Helminthosporium Leaf Spot Rad 4 Northern Leaf Blight Rad Southern Leaf Blight Rad Southern Rust Stewart's Wilt Other (Specify) Corn Lethal Necrosis	ce
4 Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) D. Ear and Kernel Rots		 	4 Anthracnose Stalk Rot Diplodia Stalk Rot Fusarium Stalk Rot Gibberella Stalk Rot Other (Specify)	
_ Aspergillus Ear and Kernel Rot (Aspergillus flavus) _ Diplodia Ear Rot (Stenocarpella maydis) 3 Fusarim Ear and Kernel Rot (Fusarium moniliforme) _ Gibberella Ear Rot (Gibberella zeae) _ Other (Specify)		 	_ Aspergillus Ear & Kernel Rot _ Diplodia Ear Rot _ Separium Ear & Kernel Rot _ Gibberella Ear Rot Other (Specify)	
Application Variety Data	Page 3	ļ	Standard Inbred Data	

Note: Use chart on first page to choose color codes for color traits.

Application Variety Data		Pag	9 4	Standard Inbred Data
11. INSECT RESISTANCE if not tested	(Rate from 1(most susceptible)	to 9 (most resistant St. Dev.); Leave blank Sample Size	l St. Dev. Sample Si
_ Banks Grass Mite Corn Earworm (Helicove	(Oligonychus pratensis)			I Banks Grass Mite I Corn Earworm
_ Leaf Feeding				I _ Leaf Feeding
	mg larval wt.	·		
_ Ear Damage	Rhopalosiphum maidis)			Ear Damage
	Rnopalosiphum maidis) Carpophilus dimidiatus)			Corn Leaf Aphid Corn Sap Beetle
European Corn Borer (C				I European Corn Borer
	oically Whorl Leaf Feeding)			I 1 st Generation
	ypically Leaf Sheath-Collar Fee	ding)		I _ 2 nd Generation
	cm tunneled/plant			
Fall Armyworm (Spodop	tera frugiperda)			I Fall Armyworm
_ Leaf-Feeding	ma lancel ut			Leaf-Feeding
Silk-Feeding Maize Weevil (Sito	, _mg larvar wt. phikus zeamais)			Maize Weevil
	princip Zournald)			_ Waize vveevii
Northern Rootworn	n (Diabrotica barberi)			I Northern Rootworm
	n (Diabrotica undecimpunctata)			Southern Rootworm
Southwestern Corn Bore	er (Diatraea grandiosella)			I Southwestern Corn Borer
_ Leaf Feeding				Leaf Feeding
Stalk Tunneling:	cm tunneled/plant			
two-spotted Spide	r Mite (Tetranychus urticae) ı (Diabrotica virgifera virgifera)			I _ Two-spotted Spider Mite I _ Western Rootworm
Other (Specify)	(Diablotica viigilela viigilela)			United the control of
12. AGRONOMIC TRAITS:				1
	days after anthesis) (Rate on sca	ale from 1=worst to	9=excellent)	I <u>3</u> Stay Green
% Dropped Ears (at _ % Pre-anthesis Bri	65 days after anthesis)			I % Dropped ears
% Pre-anthesis Ro				I % Pre-anthesis Brittle Snapping I % Pre-anthesis Root Lodging
	ot Lodging oot Lodging (at 65 days after ant	hesis)		Post-anthesis Root Lodging
	of Inbred Per Se (at 12-13% gra			5,727.0 Yield
13. MOLECULAR MARKER	S: (0=data unavailable; 1=data a	vailable but not sup	pplied; 2=data suppli	ied.)
1 Isozymes	_ RFLP's	_ RAPD's		_ Other (Specify)
REFERENCES:				
Butler, D.R. 1954. A System	for the Classification of Corn Int	ored Lines. PhD Th	esis, Ohio University	/ ·
Emerson, R.A., G.W. Beadle	e, and A.C. Fraser, 1935. A sumi	mary of Linkage Sti	idies in Maize. Corn	iell A.E.S., Mem. 180.
Farr, D.F., G.F. Bills, G.P. C. Society, St. Paul, MN.	hamuris, A.Y. Rossman. 1989. F	Fungi on Plant Prod	ucts in the United St	tates. The American Phytopathological
Jugenheimer, R.W. 1976. Co McGee, D.C. 1988. Maize Di Munsell Color Chart for Plant The Mutants of Maize. 1968. Shurtleff, M.C. 1980. Compe	: Culture, Processing, Products. prin: Improvement, Seed Product seases. APS Press, St. Paul, M : Tissues. Macbeth. P.O. Box 23 Crop Science Society of Americ ridium of Corn Diseases. APS P	ion, and Uses. Joh N. 150 pp. O. Newburgh, N.Y. a. Madison, WI. ress, St. Paul, MN.	n Wiley & Sons, Nev 12551-0230 105 pp.	omy Monograph 18. ASA, CSSA, SSSA,
Madison, WI.			ima Edition. Agrono	my monograph 10. AdA, OddA, OddA,
Stringfield, G.H. Maize Inbred U.S. Department of Agricultu	d Lines of Ohio A.E.S., Bul. 831. re 1936, 1937. Yearbook.	1959.		

COMMENTS (e.g. state how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in Exhibit D)
Insect, disease, brittle snapping and root lodging data are collected mainly from environment where variability for the trait
can be obtained within the experiment.

CLARIFICATION OF DATA IN EXHIBITS B AND C

Please note the data presented in Exhibit B and C, "Objective Description of Variety," are collected primarily at Johnston and Dallas Center, Iowa. The data in Table 1 are from two sample t-tests using data collected in Johnston and Dallas Center, IA. These traits in exhibit B collectively show distinct differences between the two varieties.

Exhibit D

Variety PHAR1 has greater Grey Leaf Spot resistance than W64A.

		•	GLFSPT score ABS	GLFSPT score ABS		GLFSPT score ABS	GLFSPT score ABS	GLFSPT score ABS
Variety1	Variety2	Year	Mean1	Mean2	#Locs	Diff	SD1	SD2
PHAR1	W64A	2002	5	2	4	3	2.3	8.0
PHAR1	W64A	2003	4.3	2.1	4	2.1	0.5	1.3
PHAR1	W64A	2004	5.4	2.1	8	3.3	1.2	0.6
PHAR1	W64A	2005	4.8	1.2	3	3.7	1.3	0.3
PHAR1	W64A	2006	4.5	2.4	4	2.1	0.6	1
PHAR1	W64A	Overall	4.9	2	23	2.9	1.2	0.9

GLFSPT = GRAY LEAF SPOT (*Cercospora zeae-maydis*). A 1 to 9 visual rating indicating the resistance to Gray Leaf Spot. A higher score indicates a higher resistance. Data are collected only when sufficient selection pressure exists in the experiment measured.

Variety PHAR1 has lower percent discard seed than PHCPR.

			KSZDCD pct ABS		KSZDCD pct ABS			KSZDCD pct ABS
Variety1	Variety2	EXPT_YEAR	Mean1	Mean2	#Locs	Diff	SD1	SD2
PHAR1	PHCPR	2002	1.3	11.0	3	-9.7	0.6	3.0
PHAR1	PHCPR	2004	2.0	6.0	4	-4.0	0.8	8.0
PHAR1	PHCPR	Overall	1.7	8.1	7	-6.4	0.8	3.2

KSZDCD = KERNEL SIZE DISCARD. The percent of discard seed; calculated as the sum of discarded tip kernels and extra large kernels.

REPRODUCE LOCALLY. Include form number and edition date on all reproductions.	FORM APPROVED - OMB NO. 058	1-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determin certificate is to be issued (7 U.S.C. 2421). confidential until the certificate is issued (7	e if a plant variety protection The information is held
1.NAME OF APPLICANT(S) PIONEER HI-BRED INTERNATIONAL, INC.	2.TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME PHAR1
4 .ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country) 7301 NW 62 nd AVENUE	5.TELEPHONE (include area code) 515-270-4051	6. FAX (include area code) 515-253-2125
P.O.BOX 85 JOHNSTON, IA 50131-0085	7.PVPO NUMBER	00500253
	9) 	
9.Is the applicant (individual or company) a U.S. national or a U.S. based company	? If no, give name of country. 🛛 YES	S NO
a. If the original rights to variety were owned by individual(s), is (are) the original rights to variety were owned by individual(s), is (are) the original rights to variety were owned by a company(ies), is (are) the		
11. Additional explanation on ownership (Trace ownership from original breeder to Pioneer Hi-Bred International, Inc. (PHI), Des Moines, Iowa, and/or its wholly is the employer of the plant breeders involved in the selection and developme Corporation has the sole rights and ownership of PHAR1pursuant to written c such variety was created. No rights to this variety are retained by any individ	owned subsidiary Pioneer Overseas Corpo ent of PHAR1. Pioneer Hi-Bred Internation contracts that assign all rights in the variety	oration (POC), Des Moines, Iowa al and/or Pioneer Overseas
PLEASE NOTE: Plant variety protection can only be afforded to the owners (not licensees) who meet the fo	ollowing criteria:	

- 1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definitions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal employment opportunity provide and employer.